

Discrete Symmetries and Transformations of the Hubbard Model.

J.P.Wallington and James F.Annett.

*University of Bristol, H.H. Wills Physics Laboratory, Royal Fort, Tyndall Ave, Bristol BS8 1TL,
United Kingdom.*

(February 1, 2008)

Abstract

We show that, in addition to $SO(4)$, the Hubbard model at half filling on a bipartite lattice has a group of discrete symmetries and transformations. A unique Hubbard-Stratonovich decomposition of the interaction term, incorporating both spin and pseudospin variables on an equal footing, is found in which these symmetries are manifestly present. The consequences of this at the mean field and one loop renormalisation group levels are discussed.

Pacs numbers: 71.10.Fd,71.27.+a

The Hubbard model is a simple, generic model of correlated electrons on a lattice. It is defined by the Hamiltonian:

$$\hat{H} = \hat{H}_0 + \hat{H}_I - \mu' \sum_{r,\sigma} n_\sigma(r), \quad (1)$$

$$\hat{H}_0 = - \sum_{\langle rr' \rangle, \sigma} t_{rr'} \left(c_\sigma^\dagger(r) c_\sigma(r') + h.c. \right), \quad (2)$$

$$\hat{H}_I = U \sum_r \left[n_\uparrow(r) - \frac{1}{2} \right] \left[n_\downarrow(r) - \frac{1}{2} \right], \quad (3)$$

where $t_{rr'}$ connects only nearest-neighbour lattice sites and $\mu' = \mu - \frac{U}{2}$ controls the number density of the system. U is the interaction energy.

The model was introduced by Hubbard [1] as a model of itinerant ferromagnets. Since then the repulsive case ($U > 0$) has been used to describe both ferro- and antiferromagnetism [2,3] and d -wave superconductivity [4,5]. For an attractive interaction ($U < 0$) it has been applied to the study of BCS s -wave superconductivity [6] and the transition to the Bose condensation of strongly bound pairs [7]. Interest in the Hubbard model on a 2D square lattice has been particularly intense since the suggestion by Anderson [8] that it is relevant to the description of the high- T_c compounds.

Despite this interest the Hubbard model remains unsolved except in 1D. However, in recent years, there have been a number of exact results proved. Yang and Zhang [9] have shown, on a bipartite lattice at half-filling and zero field, the existence of an $SO(4) \equiv SU_S(2) \otimes SU_R(2)/Z_2$ symmetry. The group's action generates rotations of the spin and ‘pseudospin’ vectors, $\mathbf{S}$ and $\mathbf{R}$:

$$\mathbf{S} \equiv \frac{1}{2} \Psi_S^\dagger(r) \boldsymbol{\sigma} \Psi_S(r), \quad \Psi_S(r) = \begin{pmatrix} c_\uparrow(r) \\ c_\downarrow(r) \end{pmatrix}, \quad (4)$$

$$\mathbf{R} \equiv \frac{1}{2} \Psi_R^\dagger(r) \boldsymbol{\sigma} \Psi_R(r), \quad \Psi_R(r) = \begin{pmatrix} c_\uparrow(r) \\ e^{i\pi \cdot r} c_\downarrow^\dagger(r) \end{pmatrix}, \quad (5)$$

where $\boldsymbol{\sigma}$ is the vector of Pauli matrices and π is the vector $(\pi, \pi, \dots)$ in d dimensions. The pseudospin vector comprises superconducting pair creation/annihilation operators and the number operator. A consequence of pseudospin rotational invariance is the existence, at

half filling, of a degenerate superconducting/charge density wave (S-CDW) ordering [10,11]. Östlund [12] has given a thorough account of the $SO(4)$ symmetry, including an enumeration of the possible mean field phases of the model.

Recently Zhang [13] proposed an approximate $SO(5)$ symmetry connecting antiferromagnetism to d -wave superconductivity. The proposal offers the possibility of understanding the phase diagram of the cuprates and has attracted much attention. The symmetry is not exact, unlike $SO(4)$, but becomes valid under the action of the renormalisation group.

In this work we examine exact symmetries. We show that in addition to the continuous $SO(4)$ there are non-trivial discrete symmetries. These play a similar role to CPT invariance in relativistic quantum mechanics. In fact, this analogy is very close, as will be clear below, because $SO(4)$ is the Euclidean analogue of the Lorentz group, $SO(3,1)$. We show that these symmetries, augmented by the addition of the Lieb-Mattis transformation, are very useful in resolving an ambiguity inherent in any field theoretic treatment based upon the Hubbard-Stratonovich transformation [14,15]:

$$e^{\frac{1}{2}\mathcal{F}^2} \equiv \int \mathcal{D}[b] e^{-\frac{1}{2}b^2 - \mathcal{F}b}, \quad (6)$$

where $\mathcal{F}$ is a bilinear Fermi operator and b is a Bose field. The use of this identity converts the problem of interacting fermions into one of free fermions interacting with bosons. To employ this technique the quartic term, $\hat{H}_I$, must be written as the square of a bilinear operator. The method is ambiguous because there are many ways (actually a continuous infinity) of doing this [16]:

$$\hat{H}_I = \sum_r \left[-\frac{U}{3} (1 - \alpha) \mathbf{S} \cdot \mathbf{S} + \frac{U}{3} (1 + \alpha) \mathbf{R} \cdot \mathbf{R} \right]. \quad (7)$$

Whilst this is formally exact for all α , each choice behaves differently under approximation schemes (e.g. mean field theory, RPA, renormalisation group). It is customary to pick a decomposition which is most convenient for the problem at hand; for instance $\alpha = -1$ for magnetism or $\alpha = +1$ for superconductivity. We will show, however, using the discrete transformation properties of the Hamiltonian, that there is a single ‘best’ decomposition.

Uniquely, approximation techniques applied to this decomposition preserve all the symmetry and transformation properties of the full Hamiltonian.

It is convenient to introduce a four spinor:

$$\Psi(r) \equiv \left(c_{\uparrow}(r), c_{\downarrow}(r), e^{i\pi \cdot r} c_{\uparrow}^{\dagger}(r), e^{i\pi \cdot r} c_{\downarrow}^{\dagger}(r) \right)^T, \quad (8)$$

in terms of which both $\mathbf{S}$ and $\mathbf{R}$ may be constructed. Equivalently a 2×2 matrix form is possible [17].

We consider three discrete symmetries; charge conjugation, $\hat{\mathcal{C}}$, parity, $\hat{\mathcal{P}}$, and time reversal, $\hat{\mathcal{T}}$, also the Lieb-Mattis canonical transformation, $\hat{Z}$.

Charge Conjugation, $\hat{\mathcal{C}}$. At half filling ($\mu' = 0$) on a bipartite lattice, for every state of energy $\epsilon(\mathbf{k})$ below the Fermi energy there is a corresponding state of energy $\epsilon(\mathbf{k} + \pi) = -\epsilon(\mathbf{k})$ above it. Under charge conjugation a hole of momentum $\mathbf{k}$ is equivalent to a particle of momentum $\mathbf{k} + \pi$. Interpreting filled states as the Dirac sea and holes as positrons, the situation is analogous to the relativistic case. With respect to the four spinor, $\Psi(r)$, charge conjugation is defined by:

$$\hat{\mathcal{C}} \equiv \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \otimes \hat{Y}, \quad (9)$$

where $\hat{Y}$ is the antilinear complex conjugation operator. Its effects on the spin and pseudospin operators are as follows:

$$\hat{\mathcal{C}} \mathbf{R} \hat{\mathcal{C}}^{-1} = -\mathbf{R}, \quad (10)$$

$$\hat{\mathcal{C}} S^{\pm} \hat{\mathcal{C}}^{-1} = S^{\pm}, \quad (11)$$

$$\hat{\mathcal{C}} S_z \hat{\mathcal{C}}^{-1} = S_z. \quad (12)$$

Parity, $\hat{\mathcal{P}}$, leaves $\mathbf{S}$, $\mathbf{R}$ and the Hamiltonian unchanged and is of little use to us here.

Time Reversal, $\hat{\mathcal{T}}$, reverses the signs of the linear and angular momenta:

$$\hat{\mathcal{T}}\hat{\mathbf{P}}\hat{\mathcal{T}}^{-1} = -\hat{\mathbf{P}}, \quad \hat{\mathcal{T}}\hat{\mathbf{S}}\hat{\mathcal{T}}^{-1} = -\hat{\mathbf{S}}. \quad (13)$$

Cooper pairs are time reversal invariant as is charge, thus:

$$\hat{\mathcal{T}}R^\pm\hat{\mathcal{T}}^{-1} = R^\pm, \quad \hat{\mathcal{T}}R_z\hat{\mathcal{T}}^{-1} = R_z. \quad (14)$$

Using these conditions, we may deduce the form of the time reversal operator acting on $\Psi(r)$:

$$\hat{\mathcal{T}} \equiv \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \otimes \hat{Y}. \quad (15)$$

The antilinear operator, $\hat{Y}$, is necessary to ensure that the commutation relations, $\hat{P}_\alpha\hat{Q}_\alpha - \hat{Q}_\alpha\hat{P}_\alpha = i\hbar$ are preserved under $\hat{\mathcal{T}}$.

The *Lieb-Mattis canonical transformation*, $\hat{Z}$, is defined by: $c_\uparrow(r) \mapsto c_\uparrow(r)$, $c_\downarrow(r) \mapsto e^{i\pi \cdot \mathbf{r}} c_\downarrow^\dagger(r)$. It acts on the four-spinor as a matrix:

$$\Psi(r) \xrightarrow{\hat{Z}} \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \Psi(r). \quad (16)$$

It maps spin into pseudospin and vice versa, leaves the non-interacting Hamiltonian, $\hat{H}_0$, unchanged but reverses the sign of the interaction term, $\hat{H}_I$:

$$\mathbf{S} \xleftrightarrow{\hat{Z}} \mathbf{R}, \quad (17)$$

$$\hat{H}_0 + \hat{H}_I \xleftrightarrow{\hat{Z}} \hat{H}_0 - \hat{H}_I. \quad (18)$$

The action of $\hat{Z}$ on the operators $\hat{\mathcal{C}}$ and $\hat{\mathcal{T}}$ is noteworthy:

$$\hat{\mathcal{C}} = \hat{Z}\hat{\mathcal{T}}\hat{Z}, \quad (19)$$

implying that the behaviour of $\mathbf{S}$ under $\hat{\mathcal{T}}$ is the same as $\mathbf{R}$ under $\hat{\mathcal{C}}$.

The Discrete Group of Transformations. Using the matrix forms of $\hat{\mathcal{C}}$, $\hat{\mathcal{T}}$ and $\hat{\mathcal{Z}}$ we see that the set, $\mathcal{G} \equiv \{\pm\mathbf{I}, \pm\hat{\mathcal{C}}, \pm\hat{\mathcal{T}}, \pm\hat{\mathcal{Z}}, \pm\hat{\mathcal{C}}\hat{\mathcal{T}}, \pm\hat{\mathcal{Z}}\hat{\mathcal{C}}, \pm\hat{\mathcal{Z}}\hat{\mathcal{T}}, \pm\hat{\mathcal{C}}\hat{\mathcal{T}}\hat{\mathcal{Z}}\}$, forms a discrete group under multiplication. It comprises 16 elements, falling into ten conjugacy classes. The character table is shown in Table I.

Each of the operators $\hat{H}_0$, $\hat{H}_I$, $\mathbf{S}$ and $\mathbf{R}$ may be classified as transforming according to a certain representation: $\hat{H}_0$ transforms as the identity representation, $\Gamma^{(1)}$, $\hat{H}_I$ transforms as $\Gamma^{(2)}$ whilst $\mathbf{S}$ and $\mathbf{R}$ transform as couplets; $\begin{pmatrix} S_x \\ R_x \end{pmatrix}$, $\begin{pmatrix} S_z \\ R_z \end{pmatrix}$ as $\Gamma^{(5)}$ and $\begin{pmatrix} S_y \\ R_y \end{pmatrix}$ as $\Gamma^{(3)} \oplus \Gamma^{(4)}$.

The space of bilinear operators is spanned by $\mathbf{S}$ and $\mathbf{R}$. Therefore the quartic operator, $\hat{H}_I$, can be decomposed into products of $\mathbf{S}$ and $\mathbf{R}$. The representations of all possible products are generated by:

$$\begin{aligned} & (\Gamma^{(3)} \oplus \Gamma^{(4)} \oplus 2\Gamma^{(5)}) \otimes (\Gamma^{(3)} \oplus \Gamma^{(4)} \oplus 2\Gamma^{(5)}) \\ &= 6\Gamma^{(1)} \oplus 6\Gamma^{(2)} \oplus 4\Gamma^{(3)} \oplus 4\Gamma^{(4)} \oplus 8\Gamma^{(5)}. \end{aligned} \quad (20)$$

Discarding all but those products which transform as $\Gamma^{(2)}$ and imposing explicit spin and pseudospin invariance we arrive at the unique decomposition:

$$\hat{H}_I = \sum_r \left[-\frac{U}{3} \mathbf{S} \cdot \mathbf{S} + \frac{U}{3} \mathbf{R} \cdot \mathbf{R} \right]. \quad (21)$$

We have constructed this decomposition (corresponding to $\alpha = 0$ in equation 7) in such a way that all its symmetries are transparent and are inherited from the transformation properties of $\mathbf{S}$ and $\mathbf{R}$. Because of this it enjoys a great advantage over the other choices of α . This becomes clear when we consider approximation schemes.

Mean Field Theory. Using equation 7 we construct a mean field theory for each α :

$$\begin{aligned} \hat{H}_I^{MF} = & \sum_r \left[-\frac{U}{3} (1 - \alpha) \{2\mathbf{S} \cdot \langle \mathbf{S} \rangle - \langle \mathbf{S} \rangle \cdot \langle \mathbf{S} \rangle\} \right. \\ & \left. + \frac{U}{3} (1 + \alpha) \{2\mathbf{R} \cdot \langle \mathbf{R} \rangle - \langle \mathbf{R} \rangle \cdot \langle \mathbf{R} \rangle\} \right]. \end{aligned} \quad (22)$$

The expectations, $\langle \dots \rangle$, are taken self consistently with respect to the mean field Hamiltonian. For all α and for $U > 0$, the solutions are either paramagnetic ($T > T_c$) or antiferromagnetic

($T < T_c$) but in both cases $\langle \mathbf{R} \rangle = 0$. Similarly for $U < 0$, there is a S-CDW phase for $T < T_c$ and $\langle \mathbf{S} \rangle = 0$.

The full theory is invariant under the combined action of $\hat{Z}$ and $U \mapsto -U$, therefore we expect a good mean field theory to share this property. For instance, the values of $\langle \mathbf{S} \rangle$ (for $U > 0$) and $\langle \mathbf{R} \rangle$ (for $U < 0$) should not change. We find that this is true for the $\alpha = 0$ theory but not for any other. This can be simply understood by considering the action of $\hat{Z}$ and $U \mapsto -U$ on equation 22. The effective couplings which the fields ‘see’, $\frac{U}{3}(1 - \alpha)$ for $\mathbf{S}$ and $\frac{U}{3}(1 + \alpha)$ for $\mathbf{R}$, are exchanged, the effect of which is to alter the expectation values $\langle \mathbf{S} \rangle$ and $\langle \mathbf{R} \rangle$. This has dramatic consequences when $\alpha = \pm 1$ in which cases the fields of interest acquire zero couplings and drop out of the Hamiltonian altogether. The $\alpha = 0$ theory avoids these problems by having the same coupling, $\frac{U}{3}$, for both $\mathbf{S}$ and $\mathbf{R}$.

It is desirable that Hartree-Fock theory be reproduced by the scalar version of equation 7:

$$\hat{H}_I = \sum_r \left[-U(1 - \alpha) S_z^2 + U(1 + \alpha) R_z^2 \right], \quad (23)$$

where $S_z = \frac{1}{2}(n_\uparrow - n_\downarrow)$ and $R_z = \frac{1}{2}(n_\uparrow + n_\downarrow - 1)$. Hartree-Fock is only recovered for the $\alpha = 0$ decomposition. For $\alpha \neq 0$ there are extra terms, $n_\sigma \langle n_\sigma \rangle$, which violate the Pauli exclusion principle. These arise because mean field theory does not respect idempotency, i.e. $n_\sigma n_\sigma = n_\sigma$ but $n_\sigma \langle n_\sigma \rangle \neq n_\sigma$. The $\alpha = 0$ theory was derived without the use of idempotency and so reproduces Hartree-Fock theory automatically.

Landau-Ginzburg Theory. Because the $\alpha = 0$ decomposition retains both $\mathbf{S}$ and $\mathbf{R}$ it has a further advantage in that, combined with the use of $\Psi(r)$, it allows us to study the renormalising effects of magnetism on superconductivity and vice versa. To see these we construct a Landau-Ginzburg (LG) functional.

Using equation (21) in the Hubbard Stratonovich transformation and then integrating out the fermionic degrees of freedom leaves an effective bosonic action in terms of the fields Φ and Δ which couple to $\mathbf{S}$ and $\mathbf{R}$. In the static limit the quadratic part reads:

$$\begin{aligned}\mathcal{S}^{(2)} &= \frac{1}{2} \sum_q \left(1 - \frac{U}{3} \chi(q)\right) \Phi(q) \cdot \Phi(-q) \\ &+ \frac{1}{2} \sum_q \left(1 + \frac{U}{3} \chi(q)\right) \Delta(q) \cdot \Delta(-q),\end{aligned}\tag{24}$$

where the susceptibility is:

$$\chi(q) = - \sum_k \frac{f(\epsilon_{\mathbf{k}+\mathbf{q}}) - f(\epsilon_{\mathbf{k}})}{\epsilon_{\mathbf{k}+\mathbf{q}} - \epsilon_{\mathbf{k}}}.\tag{25}$$

To construct the LG functional we need only long wavelength (small q) excitations about stable modes. The sign difference in the expressions for Φ and Δ means that (for $U > 0$) the stable modes are at π (antiferromagnetism) and 0 (η -pairing [18]) for Φ and Δ respectively. For $U < 0$ the stable modes of Φ and Δ are at 0 (ferromagnetism) and π (superconductivity/CDW). Taking $U > 0$ and expanding in q about the stable momenta gives the LG form:

$$\begin{aligned}\mathcal{S} &= \frac{1}{2} \sum_q \left(r_\pi + \frac{q^2}{2m_\pi}\right) \Phi(q) \cdot \Phi(-q) \\ &+ \frac{1}{2} \sum_q \left(r_0 + \frac{q^2}{2m_0}\right) \Delta(q) \cdot \Delta(-q) \\ &+ u_\pi \sum_{q_1, q_2, q_3} (\Phi_1 \cdot \Phi_2) (\Phi_3 \cdot \Phi_{-1-2-3}) \\ &+ u_0 \sum_{q_1, q_2, q_3} (\Delta_1 \cdot \Delta_2) (\Delta_3 \cdot \Delta_{-1-2-3}) \\ &+ u_x \sum_{q_1, q_2, q_3} (\Delta_1 \cdot \Delta_2) (\Phi_3 \cdot \Phi_{-1-2-3}),\end{aligned}\tag{26}$$

where $\Phi_i \equiv \Phi(q_i)$ (similarly for Δ) and:

$$r_\pi = 1 - \frac{U}{3} \chi(\pi), \quad r_0 = 1 + \frac{U}{3} \chi(0),\tag{27}$$

$$\frac{1}{m_\pi} = -\frac{U}{3} \frac{\partial^2 \chi(\pi)}{\partial q^2}, \quad \frac{1}{m_0} = \frac{U}{3} \frac{\partial^2 \chi(0)}{\partial q^2},\tag{28}$$

$$u_\pi = \frac{U^2}{72} \sum_\epsilon \left[\frac{f'(\epsilon)}{\epsilon^2} - \frac{f(\epsilon)}{\epsilon^3} + \frac{1}{2\epsilon^3} \right] N(\epsilon),\tag{29}$$

$$u_0 = \frac{U^2}{216} \sum_\epsilon f'''(\epsilon) N(\epsilon),\tag{30}$$

$$u_x = \frac{U^2}{72} \sum_\epsilon \frac{f''(\epsilon)}{\epsilon} N(\epsilon),\tag{31}$$

where $N(\epsilon)$ is the non-interacting density of states. For $U < 0$ the roles of Φ and Δ are simply reversed.

Equation 26 is the starting point for a renormalisation group (RG) analysis. We employ a one loop technique *à la* Wilson-Kogut [19,20]; i.e. successive elimination of ‘fast-modes’ (large q) followed by rescaling of fields and momenta. This process renormalises the parameters $r_{\pi,0}$ and $u_{\pi,0,x}$. The analysis shows that the cross term, u_x , is irrelevant, i.e. it flows to zero under the action of the RG. Therefore at the fixed point the spin and pseudospin are decoupled from each other. This implies that for $U > 0$ the fixed point is the usual antiferromagnetic one, whilst it is S-CDW for $U < 0$. The sole effect of u_x is to renormalise T_c . Fig. 1 shows RG flows for different initial u_x but the same temperature, $T_c(u_x = 0)$. For $u_x > 0$ the flow goes to the high temperature phase implying $T_c(u_x) < T_c(0)$. Conversely for $u_x < 0$ the ordered phase (AFM or S-CDW) is reached implying $T_c(u_x) > T_c(0)$. To conclude, T_c is depressed for positive u_x but increased for negative u_x .

In summary, we have discovered a discrete group of symmetries and transformations of the Hubbard model on a bipartite lattice at half filling and enumerated its representations. Using these we have constructed a unique Hubbard Stratonovich decomposition of the Hubbard interaction term, which treats spin and pseudospin on an equal footing. At a mean field level it is superior to other decompositions because it preserves all the transformation properties of the full theory. In addition, the unique decomposition we have derived has allowed us to construct a Landau-Ginzburg functional to examine the interplay of magnetism and superconductivity/CDW. For $U > 0$, the critical temperature for the onset of antiferromagnetism, T_c^{AFM} , is found to be renormalised by η -paired fluctuations. Reciprocally, for $U < 0$, T_c^{S-CDW} is renormalised by ferromagnetic fluctuations. Critical exponents are not affected.

This work is supported by the Engineering and Physical Sciences Research Council (EPSRC) through the provision of a studentship to J.P.W. and by the Office of Naval Research Grant No. N00014-95-1-0398 (J.F.A.).

REFERENCES

- [1] J.Hubbard, Proc. R. Soc. London Ser. A, **276**, 238 (1963).
- [2] Y.Nagaoka, Phys. Rev. **147**, 392 (1966).
- [3] J.E.Hirsch, Phys. Rev. B **31**, 4403 (1985).
- [4] C.Gross, R.Joynt and T.M.Rice, Z. Phys. B **68**, 425 (1987).
- [5] D.J.Scalapino, E.Loh and J.E.Hirsch, Phys. Rev. B **34**, 8190 (1986).
- [6] R.T.Scalettar, E.Y.Loh, J.E.Gubernatis, A.Moreo, S.R.White, D.J.Scalapino, R.L.Sugar and E.Dagotto, Phys. Rev. Lett. **62**, 1407 (1989).
- [7] R.Micnas and T.Kostyrko, *Recent Developments in High Temperature Superconductivity*, P.W.Klamut and M.Kazimierski (eds.)(Springer, 1995) p. 221, and references therein.
- [8] P.W.Anderson, Science **235**, 1196 (1987).
- [9] S.C.Zhang, Phys. Rev. Lett. **65**, 120 (1990), C.N.Yang and S.C.Zhang, Mod. Phys. Lett. B **4**, 759 (1990), S.C.Zhang, Int. J. Mod. Phys. B **55**, 120 (1991), S.C.Zhang, Phys. Rev. B **42**, 1012 (1991).
- [10] P.Miller, B.Janko and B.L.Györfy, Physica C **210**, 343 (1993).
- [11] J.P.Wallington and J.F.Annett, Phil. Mag. B **76**, 815 (1997).
- [12] S.Östlund, Phys. Rev. Lett. **69**, 1695 (1992).
- [13] S.C.Zhang, Science **275**, 1089 (1997).
- [14] J.Hubbard, Phys. Rev. Lett. **3**, 416 (1959).
- [15] R.L.Stratonovich, Dokl. Akad. Nauk SSSR **115**, 1907 (1957).
- [16] R.Schumann and E.Heiner, Phys. Lett. A **134**, 202 (1988).
- [17] H.J.Schulz, Phys. Rev. Lett. **65**, 2462 (1990).

- [18] C.N.Yang, Phys. Rev. Lett. **63**, 2144 (1989).
- [19] K.G.Wilson and J.Kogut, Phys. Rep. **12c**, 75 (1974).
- [20] M.E.Fisher, Rev. Mod. Phys. **46**, 597 (1974).

FIGURES

FIG. 1. RG flows for different initial values of u_x . For $u_x = 0$ the full lines show flows starting infinitesimally above and below T_c . The broken lines are for the same temperature but non-zero initial u_x .

TABLES

TABLE I. Character Table for the group of $\hat{\mathcal{C}}$, $\hat{\mathcal{T}}$ and $\hat{\mathcal{Z}}$.

	I	-I	$\hat{\mathcal{C}}\hat{\mathcal{T}}$	$-\hat{\mathcal{C}}\hat{\mathcal{T}}$	$\hat{\mathcal{C}}$ $\hat{\mathcal{T}}$	$-\hat{\mathcal{C}}$ $-\hat{\mathcal{T}}$	$\hat{\mathcal{Z}}$ $-\hat{\mathcal{C}}\hat{\mathcal{T}}\hat{\mathcal{Z}}$	$-\hat{\mathcal{Z}}$ $\hat{\mathcal{C}}\hat{\mathcal{T}}\hat{\mathcal{Z}}$	$\hat{\mathcal{C}}\hat{\mathcal{Z}}$ $\hat{\mathcal{T}}\hat{\mathcal{Z}}$	$-\hat{\mathcal{C}}\hat{\mathcal{Z}}$ $-\hat{\mathcal{T}}\hat{\mathcal{Z}}$
$\Gamma^{(1)}$	1	1	1	1	1	1	1	1	1	1
$\Gamma^{(2)}$	1	1	1	1	1	1	-1	-1	-1	-1
$\Gamma^{(3)}$	1	1	1	1	-1	-1	-1	-1	1	1
$\Gamma^{(4)}$	1	1	1	1	-1	-1	1	1	-1	-1
$\Gamma^{(5)}$	2	2	-2	-2	0	0	0	0	0	0
$\Gamma^{(6)}$	1	-1	-1	1	i	-i	1	-1	i	-i
$\Gamma^{(7)}$	1	-1	-1	1	i	-i	-1	1	-i	i
$\Gamma^{(8)}$	1	-1	-1	1	-i	i	-1	1	i	-i
$\Gamma^{(9)}$	1	-1	-1	1	-i	i	1	-1	-i	i
$\Gamma^{(10)}$	2	-2	2	-2	0	0	0	0	0	0

